

EXISTENCE RESULTS FOR FRACTIONAL IMPULSIVE INTEGRO-DIFFERENTIAL EQUATIONS WITH NONLOCAL CONDITION

M.H.M.RASHID ⁽¹⁾ AND A. AL-OMARI ⁽²⁾

ABSTRACT. This paper is mainly concerned with the existence of solutions of impulsive fractional abstract integro-differential equations in Banach spaces. The results are obtained by using fixed point principles. An example is provided to illustrate the theory.

1. INTRODUCTION

Fractional differential equations have recently been addressed by several researchers for a variety of problems. The interest in the study of differential equations of fractional order lies in the fact that fractional derivatives provide an excellent tool for the description of memory and hereditary properties of various materials and processes. With this advantage, the fractional-order models become more realistic and practical than the classical integer-order models, in which such effects are not taken into account. As a matter of fact, fractional differential equations arise in many

2000 *Mathematics Subject Classification.* 40H05, 46A45.

Key words and phrases. Existence of solution, Nonlinear fractional equation, Fractional calculus, Impulsive condition, Nonlocal condition.

Copyright © Deanship of Research and Graduate Studies, Yarmouk University, Irbid, Jordan.

Received: Jan 25, 2012

Accepted : Oct 10 , 2012 .

engineering and scientific disciplines such as physics, chemistry, biology, economics, control theory, signal and image processing, biophysics, blood flow phenomena, aerodynamics, fitting of experimental data, etc. [21, 28, 29, 30] . For some recent development on the topic, see [9, 22, 23] and the references therein.

In [18, 20] the authors have proved the existence of solutions of abstract differential equations by using semigroup theory and fixed point theorem. Many partial fractional differential equations can be expressed as fractional differential equations in some Banach spaces [19].

The study of nonlocal Cauchy problem for abstract evolution differential equations has been initiated by Byszewski [14]. Subsequently several authors [15, 25] have discussed the problem for different types of nonlinear differential and integrodifferential equations including functional differential equations in Banach spaces. Balachandran et al. [4, 10, 11] established the existence of solutions of quasilinear integrodifferential equations with nonlocal conditions where the quasilinear operator is unbounded. The existence of solutions of fractional abstract differential equations with nonlocal initial condition was recently investigated by NGuérékata [26] and Balachandran and Park [10]. Numerical experiments for fractional models on population dynamics are discussed in [26]. On the other hand, the theory of impulsive differential equations has undergone rapid development over the years and played a very important role in modern applied mathematical models of real processes arising in phenomena studied in physics, population dynamics, chemical technology, biotechnology and economics: see for instance the monograph by Lakshmikantham et al. [24]. Benchohra and Seba [13] studied the existence of fractional impulsive differential equations in Banach spaces while Balachandran and Kiruthika [5] discussed

the existence of nonlocal Cauchy problem for semilinear fractional evolution equations where as Chang and Nieto [16] studied the same type of problem for neutral integrodifferential inclusions via fractional operators. Belmekki et al. [7] proved the existence of periodic solutions of nonlinear fractional differential equations where as Ahmad and Nieto [1] discussed the existence results for nonlinear boundary value problem of fractional integrodifferential equations with integral boundary conditions. Recently, the study of impulsive differential equations has attracted a great deal of attention in fractional dynamics and its theory has been treated in several works [12, 24].

In this paper, motivated by [12], we prove the existence and uniqueness of solution to impulsive semi-linear fractional integro-differential equations by using the fractional calculus and fixed point theorems. This work initiates new avenues for obtaining numerical solutions of impulsive fractional integrodifferential equations.

The paper is organized as follows. In section 2 we recall the definition of the Caputo fractional derivative and Riemann-Liouville integral. In section 3 we study the existence of solution to Semilinear Evolution problem and the generalization to impulsive Evolution problem in section 4. Section 5 is devoted to study the existence of solution of the impulsive evolution equation with non-local condition. An example to illustrate our results is given in section 6.

2. PRELIMINARIES

In this section, we shall introduce some basic definitions, notations and lemmas which are used throughout this paper.

Definition 2.1. Let $f : \mathbb{R}^+ \rightarrow \mathbb{R}$ be a continuous function on \mathbb{R}^+ and $\alpha \geq 0$. Then the expression

$$I_0^\alpha f(t) = \frac{1}{\Gamma(\alpha)} \int_a^t (t-s)^{\alpha-1} ds \quad t > 0$$

is called the Riemman-Liouville integral of order α .

Definition 2.2. ([30]) Let $f : \mathbb{R}^+ \rightarrow \mathbb{R}$. The Caputo fractional derivative of order α of f is defined by

$$D_t^\alpha f(t) = \frac{1}{\Gamma(n-\alpha)} \int_0^t (t-s)^{n-\alpha-1} f^{(n)}(s) ds \quad t > 0,$$

where $\alpha \in (n-1, n)$, $n \in \mathbb{N}$.

Let $\mathbf{B}(X)$ denote the algebra of all bounded linear operator acting on a Banach space X . Now let us consider the set of functions

$$PC(I, X) = \{u : I \rightarrow X : u \in \mathcal{C}((t_{k-1}, t_k], X), k = 1, 2, \dots, m \text{ and}$$

$$\text{there exist } u(t_k^-) \text{ and } u(t_k^+), k = 1, 2, \dots, m \text{ with } u(t_k^-) = u(t_k)\}.$$

endowed with the norm

$$\|u\|_{PC} = \sup_{t \in I} \|u(t)\|$$

$(PC(I, X), \|\cdot\|_{PC})$ is a Banach space, see [24].

3. SEMILINEAR EVOLUTION PROBLEM

Consider the linear fractional impulsive evolution equation

$$(3.1) \quad \begin{aligned} D_t^\alpha u(t) &= A(t)u(t) + f(t, u(t)) \quad t \in [0, T], \quad t \neq t_k \quad 0 < \alpha \leq 1 \\ \Delta u|_{t=t_k} &= I_k(u(t_k^-)) \quad k = 1, \dots, m. \end{aligned}$$

$$u(0) = u_0 \in X$$

where $A(t)$ is a bounded linear operator on a Banach space X , $I_k : X \rightarrow X$, $k = 1, \dots, m$. and $u_0 \in X$, $0 = t_0 < t_1 < t_2 < \dots < t_m < t_{m+1} = T$. $\Delta u|_{t=t_k} = u(t_k^+) - u(t_k^-)$, $u(t_k^+) = \lim_{h \rightarrow 0^+} u(t_k + h)$ and $u(t_k^-) = \lim_{h \rightarrow 0^-} u(t_k + h)$ represent the right and left limits of $u(t)$ at $t = t_k$.

Definition 3.1. By a solution of the abstract Cauchy problem 3.1 we mean an abstract function u such that the following condition are satisfied

- (i) $u \in PC(I, X)$ and $u \in D(A(t))$ for all $t \in I' \subseteq I$;
- (ii) $\frac{\partial^\alpha u}{\partial t^\alpha}$ exists on I' where $0 < \alpha < 1$;
- (iii) u satisfies Eq. (3.1) on I' and satisfy the conditions

$$\Delta u|_{t=t_k} = I_k(u(t_k^-)) \quad k = 1, \dots, m.$$

$$u(0) = u_0 \in X$$

Now, we assume the following conditions to prove the existence of a solution of the Eq. (3.1).

(H1) $A(t)$ is a bounded linear operator on X for each $t \in I$ and the function $t \rightarrow A(t)$ is continuous in the uniform operator topology and there exists a constant M such that

$$M = \max_{t \in I} \|A(t)\|.$$

(H2) The functions $I_k : X \rightarrow X$ are continuous and there exists a constant $L_1 > 0$ and $\rho_1 > 0$ such that

$$\|I_k(u) - I_k(v)\| \leq L_1 \|u - v\| \quad \text{and} \quad \|I_k(u)\| < \rho_1 \|u\| \quad \text{for each } u, v \in X \text{ and } k = 1, 2, \dots, m,$$

and $f : I \times X \rightarrow X$ is continuous and there exists a constant L_2 such that

$$\|f(t, u) - f(t, v)\| \leq L_2 \|u - v\| \quad \text{for all } u, v \in X.$$

For brevity let us take $\frac{T^\alpha}{\Gamma(\alpha+1)} = \gamma$.

Definition 3.2. A function $u \in PC(I, X)$ solution of the fractional integral equation

$$u(t) = u_0 + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} (A(s)u(s) + f(s, u(s))) ds$$

will be called a mild solution of Problem 3.1.

Lemma 3.3. Let $f : I \times X \rightarrow X$ be a continuous function and A be a bounded linear operator. If $u \in C(I, X)$ is a mild solution of (3.1) in the sense of Definition 3.2, then for any $t \in (t_{k-1}, t_k]$, $k = 1, \dots, m$, then Eq. 3.1 is equivalent to the integral equation

$$\begin{aligned} u(t) = & u_0 + \frac{1}{\Gamma(\alpha)} \sum_{j=1}^k \int_{t_{j-1}}^{t_j} (t_j - s)^{\alpha-1} (A(s)u(s) + f(s, u(s))) ds \\ & + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} (A(s)u(s) + f(s, u(s))) ds \\ & + \sum_{0 < t_k < t} I_k(u(t_k^-)) \quad \text{if } t \in (t_k, t_{k+1}] \end{aligned} \quad (3.2)$$

Proof. Using Definition 3.2, we have

$$u(t) = u_0 + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds \quad t \in [0, t_1],$$

leads to

$$u(t_1) = u_0 + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} (t_1 - s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds$$

since $u(t_1^+) = u(t_1^-) + I_1(u(t_1^-)) = u(t_1) + I_1(u(t_1^-))$, then

$$u(t_1^+) = u_0 + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} (t_1 - s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds + I_1(u(t_1^-))$$

Moreover, for $t \in (t_1, t_2]$,

$$\begin{aligned} u(t) &= u(t_1^+) + \frac{1}{\Gamma(\alpha)} \int_{t_1}^t (t-s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds \\ &= u_0 + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} (t-s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_1}^t (t-s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds + I_1(u(t_1^-)) \end{aligned}$$

For $t = t_2$ and $t = t_2^+$,

$$\begin{aligned} u(t_2) &= u_0 + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} (t_1-s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_1}^{t_2} (t_2-s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds + I_1(u(t_1^-)) \end{aligned}$$

since $u(t_2^+) = u(t_2^-) + I_2(u(t_2^-)) = u(t_2) + I_2(u(t_2^-))$, then

$$\begin{aligned} u(t_2^+) &= u_0 + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} (t_1-s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_1}^{t_2} (t_2-s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds \\ &\quad + I_1(u(t_1^-)) + I_2(u(t_2^-)) \end{aligned}$$

Hence for any $t \in (t_2, t_3]$,

$$\begin{aligned} u(t) &= u(t_2^+) + \frac{1}{\Gamma(\alpha)} \int_{t_2}^t (t-s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds \\ &= T(t)u_0 + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} (t_1-s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_1}^{t_2} (t_2-s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_2}^t (t-s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds \\ &\quad + I_1(u(t_1^-)) + I_2(u(t_2^-)) \end{aligned}$$

By repeating the same procedure, we can easily deduce that

$$\begin{aligned} u(t) &= T(t)u_0 + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} [A(s)u(s) + f(s, u(s))] ds \\ &\quad + \sum_{0 < t_k < t} I_k(u(t_k^-)) \end{aligned}$$

holds for any $t \in (t_{k-1}, t_k]$, $k = 1, \dots, m$. □

Theorem 3.4. *If the hypotheses (H1) and (H2) are satisfied, then Equation 3.1 has a unique solution on I .*

Proof. The proof is based on the application of Picards iteration method. Let $M = \max_{0 \leq t \leq T} \|A(t)\|$ and define a mapping $F : PC([0, T] : X) \rightarrow PC([0, T] : X)$ by

$$\begin{aligned} Fu(t) &= u_0 + \frac{1}{\Gamma(\alpha)} \sum_{j=1}^k \int_{t_{j-1}}^{t_j} (t_j - s)^{\alpha-1} (A(s)u(s) + f(s, u(s))) ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} (A(s)u(s) + f(s, u(s))) ds \\ &\quad + \sum_{0 < t_k < t} I_k(u(t_k^-)) \quad \text{if } t \in (t_k, t_{k+1}] \end{aligned} \quad (3.3)$$

Let $u, v \in PC(I, X)$. Then from Eq. (3.3), we have for each $t \in I$

$$\|Fu(t) - Fv(t)\| \leq \frac{T^\alpha(m+1)}{\Gamma(\alpha+1)} (M + L_2) \|u - v\| + mL_1 \|u - v\|.$$

Then by induction we have

$$\|F^n u(t) - F^n v(t)\| \leq \frac{(\gamma(m+1)(M + L_2) + mL_1)^n}{n!} \|u - v\|.$$

Since $\frac{(\gamma(m+1)(M+L_2)+mL_1)^n}{n!} < 1$ for large n , then by the well-known generalization of the Banach contraction principle, F has a unique fixed point $u \in PC([0, T] : X)$. This fixed point is the unique solution of Eq.(3.1). \square

4. IMPULSIVE SEMILINEAR CAUCHY PROBLEM

Let $A(t)$ is a bounded linear operator and the nonlinear functions $f, g : I \times X \times X \rightarrow X$ and $h : D \times X \rightarrow X$ are continuous. Here $D = \{(t, s) \in \mathbb{R}^2 : 0 \leq s \leq t \leq T\}$. For brevity let

$$Hu(t) = \int_0^t h(t, s, u(s)) ds.$$

(H3) h is continuous function and there exist positive constants α_1, α_2 and α such that

$$\|h(t, s, u) - h(t, s, v)\| \leq \alpha_1 \|u - v\|$$

for all $u, v \in \mathcal{C}(I, X)$, $\alpha_2 = \max_{(t,s) \in D} \|h(t, s, 0)\|$, and $\alpha = \max\{\alpha_1, \alpha_2\}$.

Lemma 4.1. *If (H3) is satisfied, then the estimate*

$$\|Hu(t)\| \leq t(\alpha_1 \|u\| + \alpha_2)$$

$$\|Hv(t) - Hw(t)\| \leq \alpha_1 t \|v - w\|.$$

is satisfied for any $t \in I$, and $u, v, w \in \mathcal{C}(I, X)$.

We need the following additional assumptions to prove the existence of solution of the Eq. (3.1).

(H4) f, g are continuous and there exist a positive constants L, L_2, L_3, K, K_2, K_3 and N such that

$$\begin{aligned} \|f(t, u, Hu(t)) - f(t, v(t), Hv(t))\| &< L_2 \|u - v\| + L_3 \|Hu - Hv\| \\ &< (L_2 + L_3 \alpha_1 t) \|u - v\| < L \|u - v\| \quad \text{for all } u, v \in X \end{aligned}$$

$$\begin{aligned} \|g(t, u, Hu(t)) - g(t, v(t), Hv(t))\| &< K_2 \|u - v\| + K_3 \|Hu - Hv\| \\ &< (K_2 + K_3 \alpha_1 t) \|u - v\| < K \|u - v\| \quad \text{for all } u, v \in X \end{aligned}$$

$$\text{and } N = \max_{t \in I} \|f(t, 0, 0)\|, \lambda = \max_{t \in I} \|g(t, 0, 0)\|$$

(H4*) The function $q : I \rightarrow X$ is continuous and there exist constant q_t such that

$$\int_0^t |q(t-s)| ds = q_t.$$

Now consider the semilinear fractional impulsive evolution equation

$$u^{(\alpha)} = A(t)u(t) + f(t, u(t), Hu(t)) + \int_0^t q(t-s)g(s, u(s), Hu(s)) ds,$$

$$t \in [0, T], \quad 0 < \alpha \leq 1$$

$$(4.1) \quad \Delta u|_{t=t_k} = I_k(u(t_k^-)) \quad k = 1, \dots, m.$$

$$u(0) = u_0 \in X$$

The Eq. (4.1) is equivalent to the integral equation

$$\begin{aligned} u(t) = u_0 &+ \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} [A(s)u(s) + f(s, u(s), Hu(s)) \\ &+ \int_0^s q(s-\tau)g(\tau, u(\tau), Hu(\tau)) d\tau] ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t-s)^{\alpha-1} [A(s)u(s) + f(s, u(s), Hu(s)) \\ &+ \int_0^s q(s-\tau)g(\tau, u(\tau), Hu(\tau)) d\tau] ds + \sum_{0 < t_k < t} I_k(u(t_k^-)) \end{aligned}$$

Theorem 4.2. *If the hypotheses $H1$, $H2$, $H3$ and $H4$ are satisfied and if*

$$[\gamma(m+1)(M+L+q_tK)+m\rho_1] < \frac{1}{2}$$

then the fractional impulsive Eq.(4.1) has a unique solution on I .

Proof. Let $Z = PC(I, X)$. Define the mapping $\Phi : Z \rightarrow Z$ by

$$\begin{aligned} \Phi u(t) = & u_0 + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} [A(s)u(s) + f(s, u(s), Hu(s))] \\ & + \int_0^s q(s - \tau)g(\tau, u(\tau)) d\tau] ds \\ & + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} [A(s)u(s) + f(s, u(s), Hu(s))] \\ & + \int_0^s q(s - \tau)g(\tau, u(\tau), Hu(\tau)) d\tau] ds + \sum_{0 < t_k < t} I_k(u(t_k^-)) \end{aligned}$$

and we have to show that Φ has a fixed point. This fixed point is the solution of Eq. (4.1). Choose $r \geq 2[\|u_0\| + \gamma(m+1)(N+q_t\lambda)]$.

Then we can show that $\Phi B_r \subseteq B_r$, where $B_r := \{u \in Z : \|u\| \leq r\}$. From the

assumption we have

$$\begin{aligned}
\|\Phi u(t)\| &\leq \|u_0\| + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} [\|A(s)\| \|u(s)\| + \|f(s, u(s), Hu(s))\| \\
&\quad + \left\| \int_0^s q(s - \tau) g(\tau, u(\tau), Hu(\tau)) d\tau \right\|] ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} [\|A(s)\| \|u(s)\| + \|f(s, u(s), Hu(s))\| \\
&\quad + \left\| \int_0^s q(s - \tau) g(\tau, u(\tau), Hu(\tau)) d\tau \right\|] ds + \sum_{0 < t_k < t} \|I_k(u(t_k^-))\| \\
&\leq \|u_0\| + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} [\|A(s)\| \|u(s)\| \\
&\quad + \|f(s, u(s), Hu(s)) - f(s, 0, 0)\| + \|f(s, 0, 0)\| \\
&\quad + \left\| \int_0^s q(s - \tau) [g(\tau, u(\tau), Hu(\tau)) - g(\tau, 0, 0)] d\tau \right\| \\
&\quad + \left\| \int_0^s q(s - \tau) g(\tau, 0, 0) d\tau \right\|] ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} [\|A(s)\| \|u(s)\| \\
&\quad + \|f(s, u(s), Hu(s)) - f(s, 0, 0)\| + \|f(s, 0, 0)\| \\
&\quad + \left\| \int_0^s q(s - \tau) [g(\tau, u(\tau), Hu(\tau)) - g(\tau, 0, 0)] d\tau \right\| \\
&\quad + \left\| \int_0^s q(s - \tau) g(\tau, 0, 0) d\tau \right\|] ds + \sum_{0 < t_k < t} \|I_k(u(t_k^-))\| \\
&\leq \|u_0\| + \gamma \|u\| [(m+1)[M+L+q_t K] + \rho_1 m] + \gamma(m+1)(N+q_t \lambda) \\
&\leq \|u_0\| + \gamma(m+1)(N+q_t \lambda) + r [\gamma(m+1)(M+L+q_t K) + m \rho_1] \\
&\leq r
\end{aligned}$$

Thus Φ maps B_r into itself. Now, for $u_1, u_2 \in Z$, we have

$$\begin{aligned}
\|\Phi u_1(t) - \Phi u_2(t)\| &\leq \|u_0\| + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} [\|A(s)(u_1(s) - u_2(s))\| \\
&\quad + \|f(s, u_1(s), Hu_1(s)) - f(s, u_2(s), Hu_2(s))\| \\
&\quad + \left\| \int_0^s q(s - \tau) [g(\tau, u_1(\tau), Hu_1(\tau)) - g(\tau, u_2(\tau), Hu_2(\tau))] d\tau \right\|] ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} [\|A(s)(u_1(s) - u_2(s))\| \\
&\quad + \|f(s, u_1(s), Hu_1(s)) - f(s, u_2(s), Hu_2(s))\| \\
&\quad + \left\| \int_0^s q(s - \tau) [g(\tau, u_1(\tau), Hu_1(\tau)) - g(\tau, u_2(\tau), Hu_2(\tau))] d\tau \right\|] ds \\
&\quad + \sum_{0 < t_k < t} \|I_k(u_1(t_k^-)) - I_k(u_2(t_k^-))\| \\
&\leq \left[\frac{T^\alpha}{\Gamma(\alpha + 1)} (m + 1) [M + L + q_t K] + mL_1 \right] \|u_1 - u_2\| \\
&\leq [\gamma(m + 1) [M + L + q_t K] + mL_1] \|u_1 - u_2\| \\
&\leq \frac{1}{2} \|u_1 - u_2\|.
\end{aligned}$$

□

Hence Φ is a contraction mapping and therefore there exists a unique fixed point $u \in B_r$ such that $\Phi u(t) = u(t)$. Any fixed point of Φ is the solution of Equation (4.1).

5. NONLOCAL IMPULSIVE SEMILINEAR CAUCHY PROBLEM

In this section we discuss the existence of solution of the impulsive evolution equation (4.1) with nonlocal condition of the form

$$u(0) + \chi(u) = u_0 \tag{5.1}$$

where $\chi : PC(I, X) \rightarrow X$ is a given function which satisfies the following condition

(H5) $\chi : PC(I, X) \rightarrow X$ is continuous and there exists a constant $G > 0$ such that

$$\|\chi(u) - \chi(v)\| \leq G \|u - v\|_{PC} \quad \text{for } u, v \in PC(I, X).$$

Theorem 5.1. *If the hypotheses (H1), (H2), (H3), (H4) and (H5) are satisfied and if*

$$[G + \gamma [(m+1)(M + L + q_t K)] + mL_1] < \frac{1}{2}$$

then the fractional impulsive evolution Equation (4.1) with nonlocal condition (5.1) has a unique solution on I .

Proof. We want to prove that the operator $\Psi : Z \rightarrow Z$ defined by

$$\begin{aligned} \Psi u(t) = & u_0 - \chi(u) + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} A(s) u(s) ds \\ & + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} A(s) u(s) ds \\ & + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} \left[f(s, u(s), Hu(s)) + \int_0^s q(s - \tau) g(\tau, u(\tau), Hu(\tau)) d\tau \right] ds \\ & + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} \left[f(s, u(s), Hu(s)) + \int_0^s q(s - \tau) g(\tau, u(\tau), Hu(\tau)) d\tau \right] ds \\ & + \sum_{0 < t_k < t} I_k(u(t_k^-)) \end{aligned}$$

has a fixed point. This fixed point is then a solution of Equation (4.1) with nonlocal condition (5.1). Choose $r \geq 2(\|u_0\| + \|\chi(0)\| + (m+1)N\gamma)$. Then it is easy to see that $\Psi B_r \subset B_r$.

Further, for $u, v \in Z$, we have

$$\begin{aligned}
\|\Psi u - \Psi v\| &\leq G \|u(t) - v(t)\| + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} \|A(s)\| \|u(s) - v(s)\| ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t_k - s)^{\alpha-1} \|A(s)\| \|u(s) - v(s)\| ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} [\|f(s, u(s), Hu(s)) - f(s, v(s), Hv(s))\| \\
&\quad + \int_0^s \|q(s - \tau)\| \|g(\tau, u(\tau), Hu(\tau)) - g(\tau, v(\tau), Hv(\tau))\| d\tau] ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} [\|f(s, u(s), Hu(s)) - f(s, v(s), Hv(s))\| \\
&\quad + \int_0^s \|q(s - \tau)\| \|g(\tau, u(\tau), Hu(\tau)) - g(\tau, v(\tau), Hv(\tau))\| d\tau] ds \\
&\quad + \sum_{0 < t_k < t} \|I_k(u(t_k^-) - I_k(v(t_k^-))\| \\
&\leq \left[G + \frac{T^\alpha}{\Gamma(\alpha + 1)} [(m + 1)(M + L + q_t K)] + mL_1 \right] \|u - v\| \\
&\leq [G + \gamma [(m + 1)(M + L + q_t K)] + mL_1] \|u - v\| \\
&\leq \frac{1}{2} \|u - v\|.
\end{aligned}$$

The result follows by the application of contraction mapping principle. \square

Our next result is based on the following well-known fixed point theorem.

Krasnoselskii Theorem. ([21]) *Let S be closed convex nonempty subset of a Banach space X . Let P, Q be two operators such that*

- (i) $Px + Qx \in S$ whenever $x, y \in S$;
- (ii) P is a contraction mapping;
- (iii) Q is compact and continuous.

Then there exists $z \in S$ such that $z = Pz + Qz$.

Now, we assume the following condition with (H4) and apply the above fixed point theorem.

(H6) $f : I \times X \times X \rightarrow X$ is continuous and there exists a continuous function $\mu \in L^1(I)$ such that

$$\|f(t, u, Hu)\| \leq \mu(t), \quad \text{for all } (t, u, Hu) \in I \times X \times X.$$

(H7) $g : I \times X \times X \rightarrow X$ is continuous and there exists a continuous function $\mu \in L^1(I)$ such that

$$\|g(t, u, Hu)\| \leq \rho(t), \quad \text{for all } (t, u, Hu) \in I \times X \times X.$$

Theorem 5.2. *Assume that (H1), (H2), (H3), (H4), (H5), (H6), and (H7) hold. If $G + \gamma(m+1)M + mL_1 < 1$, then the fractional evolution Equation (4.1) with nonlocal condition (5.1) has a solution on I .*

Proof. Choose

$$r \geq \frac{\|u_0\| + \|\chi(0)\| + \gamma(m+1)(\mu_0 + q_t\rho_0)}{1 - (G + [\gamma(m+1)M] + m\rho_1)}$$

where $\mu_0 = \sup_{t \in I} \mu(t)$, $\rho_0 = \sup_{t \in I} \rho(t)$ and define the operators P and Q on B_r as

$$\begin{aligned} Pu(t) &= u_0 - \chi(u) + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} A(s) u(s) ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} A(s) u(s) ds + \sum_{0 < t_k < t} I_k(u(t_k^-)) \end{aligned}$$

and

$$\begin{aligned} Qu(t) &= \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} \left[f(s, u(s), Hu(s)) + \int_0^s q(s - \tau) g(\tau, u(\tau), Hu(\tau)) d\tau \right] ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} \left[f(s, u(s), Hu(s)) + \int_0^s q(s - \tau) g(\tau, u(\tau), Hu(\tau)) d\tau \right] ds. \end{aligned}$$

For any $u, v \in B_r$, we have

$$\begin{aligned} \|Pu(t) + Qv(t)\| &\leq \|u_0\| + \|\chi(u) - \chi(0)\| + \|\chi(0)\| \\ &\quad + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} \|A(s)\| \|u(s)\| ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} \|A(s)\| \|u(s)\| ds + \sum_{0 < t_k < t} \|I_k(u(t_k^-))\| \\ &\quad + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} \left[\|f(s, u(s), Hu(s))\| + \int_0^s \|q(s - \tau)\| \|g(\tau, u(\tau), Hu(\tau))\| d\tau \right] ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} \left[\|f(s, u(s), Hu(s))\| + \int_0^s \|q(s - \tau)\| \|g(\tau, u(\tau), Hu(\tau))\| d\tau \right] ds \\ &\leq \|u_0\| + \|\chi(0)\| + r[G + [\gamma(m+1)M] + m\rho_1] + \gamma(m+1)(\mu_0 + q_t\rho_0) \\ &\leq r. \end{aligned}$$

Hence, we deduce that $\|Pu + Qv\| \leq r$.

Next, for any $t \in I$, $u, v \in X$ we have

$$\begin{aligned} \|Pu - Pv\| &\leq \|\chi(u) - \chi(v)\| + \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} \|A(s)\| \|u(s) - v(s)\| ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} \|A(s)\| \|u(s) - v(s)\| ds + \sum_{0 < t_k < t} \|I_k(u(t_k^-)) - I_k(v(t_k^-))\| \\ &\leq G \|u - v\| + \gamma(m+1)M \|u - v\| + mL_1 \|u - v\| \\ &\leq [G + \gamma(m+1)M + mL_1] \|u - v\|. \end{aligned}$$

And since $G + \gamma(m+1)M + mL_1 < 1$, then P is a contraction mapping.

Now, let us prove that Q is continuous and compact.

Let $\{u_n\}$ be a sequence in B_r , such that $u_n \rightarrow u$ in B_r . Then $f(s, u_n(s), Hu_n(s)) \rightarrow f(s, u(s), Hu(s))$, and $g(s, u_n(s), Hu_n(s)) \rightarrow g(s, u(s), Hu(s))$ $n \rightarrow \infty$ because the functions f and g are continuous on $I \times X \times X$. Now, for each $t \in I$, we have

$$\begin{aligned} \|Qu_n - Qu\| &\leq \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} [\|f(s, u_n(s), Hu_n(s)) - f(s, u(s), Hu(s))\| \\ &\quad + \int_0^s \|q(s - \tau)\| \|g(\tau, u_n(\tau), Hu_n(\tau)) - g(\tau, u(\tau), Hu(\tau))\| d\tau] ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} [\|f(s, u_n(s), Hu_n(s)) - f(s, u(s), Hu(s))\| \\ &\quad + \int_0^s \|q(s - \tau)\| \|g(\tau, u_n(\tau), Hu_n(\tau)) - g(\tau, u(\tau), Hu(\tau))\| d\tau] ds \end{aligned}$$

Consequently, $\lim_{n \rightarrow \infty} \|Qu_n(t) - Qu(t)\| = 0$. In other word, Q is continuous.

Let's now note that Q is uniformly bounded on B_r . This follows from the inequality

$$\begin{aligned} \|Qu(t)\| &\leq \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} \left[\|f(s, u(s), Hu(s))\| + \int_0^s \|q(s - \tau)\| \|g(\tau, u(\tau), Hu(\tau))\| d\tau \right] ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_k}^t (t - s)^{\alpha-1} \left[\|f(s, u(s), Hu(s))\| + \int_0^s \|q(s - \tau)\| \|g(\tau, u(\tau), Hu(\tau))\| d\tau \right] ds \\ &\leq \gamma[(m+1)(\mu_0 + q_t \rho_0)]. \end{aligned}$$

*/ Now, lets prove that $Qu, u \in B_r$ is equicontinuous. Let $t_1, t_2 \in I, t_1 < t_2$, and let B_r be a bounded set in X . Let $u \in B_r$, we have

$$\begin{aligned}
\|Qu(t_2) - Qu(t_1)\| &\leq + \frac{1}{\Gamma(\alpha)} \int_{t_1}^{t_2} (t_2 - s)^{\alpha-1} [\|f(s, u(s), Hu(s))\| \\
&\quad + \int_0^s \|q(s - \tau)\| \|g(\tau, u(\tau), Hu(\tau))\| d\tau] ds \\
&= \frac{1}{\Gamma(\alpha)} \sum_{0 < t_k < t_2} \int_{t_{k-1}}^{t_k} (t_k - s)^{\alpha-1} [\|f(s, u(s), Hu(s))\| \\
&\quad + \int_0^s \|q(s - \tau)\| \|g(\tau, u(\tau), Hu(\tau))\| d\tau] ds \\
&\quad - \frac{1}{\Gamma(\alpha)} \int_0^{t_1} (t_2 - s)^{\alpha-1} [\|f(s, u(s), Hu(s))\| \\
&\quad + \int_0^s \|q(s - \tau)\| \|g(\tau, u(\tau), Hu(\tau))\| d\tau] ds \\
&\leq \frac{1}{\Gamma(\alpha)} \int_0^{t_1} [(t_2 - s)^{\alpha-1} - (t_1 - s)^{\alpha-1}] [\|f(s, u(s), Hu(s))\| \\
&\quad + \int_0^s \|q(s - \tau)\| \|g(\tau, u(\tau), Hu(\tau))\| d\tau] ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_{t_1}^{t_2} (t_2 - s)^{\alpha-1} [\|f(s, u(s), Hu(s))\| \\
&\quad + \int_0^s \|q(s - \tau)\| \|g(\tau, u(\tau), Hu(\tau))\| d\tau] ds \\
&\leq \frac{\mu_0 + q_t \rho_0}{\Gamma(\alpha + 1)} [2(t_2 - t_1)^\alpha - t_1^\alpha + t_2^\alpha]
\end{aligned}$$

As $t_2 \rightarrow t_1$, the right hand side of the above inequality tends to zero. In short we have proven that $Q(B_r)$ is relatively compact for $t \in I$. By Arzela Ascoli's theorem, Q is compact. Hence by the Krasnoselskii theorem there exists a solution of the problem (4.1) with nonlocal condition (5.1). \square

6. EXAMPLE

In this section we provide with an example to illustrate our main result. Consider the following fractional integrodifferential equation with impulsive of the form.

$$\begin{aligned}
{}^c D^\alpha u(t) &= \frac{1}{100} e^{-t} |u(t)| + f(t, u(t), Hu(t)) + \int_0^t q(t-s) g(s, u(s) Hu(s)) ds \\
t \in I = [0, 1], \quad x &\in (0, \pi) \quad t \neq \frac{1}{2}, \quad 0 < \alpha \leq 1. \\
\Delta u|_{t=\frac{1}{2}} &= \frac{|u(\frac{1}{2}^-)|}{9 + |u(\frac{1}{2}^-)|} \\
u(0) &= u_0,
\end{aligned}$$

where

$$A(t) = \frac{1}{100} e^{-t} I, \quad Hu(t) = \int_0^t e^{-\frac{1}{5}u(s)} ds$$

$$f(t, u, Hu) = \frac{e^{-t}|u|}{(49 + e^t)(1 + |u|)} + \frac{1}{10} Hu(t), \quad g(t, u, Hu) = \frac{|u|}{(t+5)^2(1 + |u|)} + \frac{1}{20} Hu(t)$$

$$q(t-s) = e^{t-s} \quad \text{and} \quad I_k(u) = \frac{|u|}{9 + |u|}, \quad u \in X$$

Let $u, v \in \mathcal{C}(I, X)$ and $t \in I$. Then we have

$$\|Hu - Hv\| = \left| \int_0^t e^{-\frac{1}{5}u(s)} ds - \int_0^t e^{-\frac{1}{5}v(s)} ds \right| \leq \frac{1}{5} \|u - v\|$$

$$\begin{aligned}
\|f(t, u, Hu) - f(t, v, Hv)\| &= \frac{e^{-t}}{49 + e^t} \left| \frac{u}{1+u} - \frac{v}{1+v} \right| + \frac{1}{10} \|Hu - Hv\| \\
&= \frac{e^{-t}|u-v|}{(49 + e^t)(1+u)(1+v)} + \frac{1}{10} \|Hu - Hv\| \\
&\leq \frac{e^{-t}}{49 + e^t} |u-v| + \frac{1}{10} \|Hu - Hv\| \\
&\leq \frac{1}{50} [|u-v| + \|Hu - Hv\|] \leq \frac{2}{50} \|u-v\|.
\end{aligned}$$

$$\begin{aligned}
\|g(t, u, Hu) - g(t, v, Hv)\| &= \frac{1}{(t+5)^2} \left| \frac{u}{1+u} - \frac{v}{1+v} \right| + \frac{1}{20} \|Hu - Hv\| \\
&= \frac{|u-v|}{(t+5)^2(1+u)(1+v)} + \frac{1}{20} \|Hu - Hv\| \\
&\leq \frac{1}{(t+5)^2} |u-v| + \frac{1}{20} \|Hu - Hv\| \\
&\leq \frac{1}{25} |u-v| + \frac{1}{20} \|Hu - Hv\| \leq \frac{5}{100} \|u-v\|.
\end{aligned}$$

Hence the condition (H1), (H4), and (H4*) hold with $M = \frac{1}{100}$, $L = \frac{2}{50}$, $K = \frac{5}{100}$, $q_t = 2$. Let $u, v \in X$. Then we have by (H2)

$$\|I_k(u) - I_k(v)\| = \left| \frac{u}{9+u} - \frac{v}{9+v} \right| = \frac{9|u-v|}{(9+u)(9+v)} \leq \frac{1}{9} |u-v|$$

Note that $L_1 = \frac{1}{9}$. Choose $r = 1, m = 1$ we shall check the condition

$$[\gamma(m+1)(M + L + q_t K) + m\rho_1] < \frac{1}{2}$$

is satisfied. Indeed

$$[\gamma(m+1)(M + L + q_t K) + m\rho_1] < \frac{1}{2} \Leftrightarrow \Gamma(\alpha+1) > \frac{75}{100}, \quad (6.1)$$

which is satisfied for some $0 < \alpha \leq 1$. Then by Theorem 4.2 the problem has a unique solution on $[0, 1]$ for some values of α satisfied condition (6.1).

Acknowledgement

The authors would like to thank the editor and the referees for useful comments and suggestions that will improve the quality of this paper.

REFERENCES

- [1] B. Ahmad , J.J. Nieto, Existence results for nonlinear boundary value problems of fractional integrodifferential equations with integral boundary conditions, *Bound. Value Probl.* Article ID. 708576 (2009), 1-11.
- [2] B. Ahmad , J.J. Nieto, A. Alsaedi, Existence and uniqueness of solutions for nonlinear fractional differential equations with non-separated type integral boundary conditions, *Acta Math. Sci.* **31** (2011), 2122–2130.
- [3] B.Ahmad, J.J. Nieto, A. Alsaedi, M. El-Shahed, A study of nonlinear Langevin equation involving two fractional orders in different intervals, *Nonlinear Anal. Real World Appl.* **13** (2012), 599–606.
- [4] K. Balachandran, K. Uchiyama, Existence of solutions of quasilinear integrodifferential equations with nonlocal condition, *Tokyo J. Math.* **23** (2000), 203-210.
- [5] K. Balachandran, S. Kiruthika, Existence of solutions of abstract fractional impulsive semi-linear evolution equations, *Electron. J. Qual. Theory Differ. Equ.* **4** (2010), 1-12.
- [6] K. Balachandran, S. Kiruthika, J.J. Trujillo, Remark on the existence results for fractional impulsive integrodifferential equations in Banach spaces, *Commun. Nonlinear Sci. Numer. Simul.* **17** (2012), 2244-2247
- [7] K. Balachandran, S. Kiruthika, J. J. Trujillo, Existence of periodic solutions for a nonlinear fractional differential equation, *Bound. Value Probl.* Article ID. 324561 (2009), 1-18.
- [8] M. Benchohra, B.A. Slimani, Existence and uniqueness of solutions to impulsive fractional differential equations, *Electron. J. Differential Equations* **10** (2009), 1–11.

- [9] K. Balachandran, J.J. Trujillo, The nonlocal Cauchy problem for nonlinear fractional integrodifferential equations in Banach spaces, *Nonlinear Anal.* **72** (2010), 4587-4593.
- [10] K. Balachandran, D.G. Park, Existence of solutions of quasilinear integrodifferential evolution equations in Banach spaces, *Bull. Korean Math. Soc.* **46** (2009), 691-700.
- [11] K. Balachandran, F. Paul Samuel, Existence of solutions for quasilinear delay integrodifferential equations with nonlocal conditions, *Electron. J. Differential Equations* **6** (2009), 1-7.
- [12] M. Benchohra, B.A. Slimani, Existence and uniqueness of solutions to impulsive fractional differential equations, *Electron. J. Differential Equations* **10** (2009), 1-11.
- [13] M. Benchohra, D. Seba, Impulsive fractional differential equations in Banach Spaces, *Electron. J. Qual. Theory Differ. Equ.* **8** (2009), 1-14.
- [14] L. Byszewski, Theorems about the existence and uniqueness of solutions of a semilinear evolution nonlocal Cauchy problem, *J. Math. Anal. Appl.* **162** (1991), 494-505.
- [15] L. Byszewski, H. Akca, Existence of solutions of a semilinear functional differential evolution nonlocal problem, *Nonlinear Anal.* **34** (1998), 65-72.
- [16] Y.K. Chang, J.J. Nieto, Existence of solutions for impulsive neutral integrodifferential inclusions with nonlocal initial conditions via fractional operators, *Numer. Funct. Anal. Optim.* **30** (2009), 227-44.
- [17] F. Chen, J.J. Nieto, Y. Zhou, Global attractivity for nonlinear fractional differential equations, *Nonlinear Anal. Real World Appl.* **13** (2012), 287-298.
- [18] M. M. El-Borai, Semigroups and some nonlinear fractional differential equations, *Appl. Math. Comput.* **149** (2004) 823-831.
- [19] M. A. El-Sayed, Fractional order diffusion wave equation, *Internat. J. Theoret. Phys.* **35** (1996), 311-322.
- [20] O.K. Jaradat, A. Al-Omari, S. Momani, Existence of the mild solution for fractional semilinear initial value problems, *Nonlinear Anal.* **69** (2008), 3153-3159.
- [21] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, in: *North-Holland Mathematics Studies*, vol. 204, Elsevier Science B.V., Amsterdam, 2006.
- [22] V. Lakshmikantham, A.S. Vatsala, Basic theory of fractional differential equations, *Nonlinear Anal.* **69** (8) (2008), 2677-2682.

- [23] V. Lakshmikantham, S. Leela, J. Vasundhara Devi, *Theory of Fractional Dynamic Systems*, Cambridge Academic Publishers, Cambridge, 2009.
- [24] V. Lakshmikantham, D.D. Bainov, P.S. Simeonov, *Theory of Impulsive Differential Equations*, Singapore, World Scientific; 1989.
- [25] J.Liang, J.Liu, T.J. Xiao, Nonlocal Cauchy problems governed by compact operator families, *Nonlinear Anal.* **57** (2004), 183-9.
- [26] NGuérékata G. M., A Cauchy problem for some fractional abstract differential equation with non local conditions, *Nonlinear Anal.* **70** (2009), 1873-6.
- [27] A. Pazy, *Semigroup of Linear Operators and Applications to Partial Differential Equations*, Springer-Verlag, New York, 1983.
- [28] I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [29] J. Sabatier, O.P. Agrawal, J.A.T. Machado (Eds.), *Advances in Fractional Calculus: Theoretical Developments and Applications in Physics and Engineering*, Springer, Dordrecht, 2007.
- [30] S.G. Samko, A.A. Kilbas, O.I. Marichev, *Fractional Integrals and Derivatives, Theory and Applications*, Gordon and Breach, Yverdon, 1993.
- [31] S. Sivasankaran, M. Mallika Arjunan, V. Vijayakumar, *Existence of global solutions for second order impulsive abstract partial differential equations*, *Nonlinear Anal.* **74** (2011), 6747–6757.

(1)DEPARTMENT OF MATHEMATICS& STATISTICS, FACULTY OF SCIENCE P.O.Box(7),MU'TAH UNIVERSITY

E-mail address: malik.okasha@yahoo.com

(2) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, AL AL-BAYT UNIVERSITY

E-mail address: alomarimutah1@yahoo.com